

FUNDAMENTAL GROUP OF SIMPLE C^* -ALGEBRAS WITH UNIQUE TRACE II

NORIO NAWATA AND YASUO WATATANI

ABSTRACT. We show that any countable subgroup of the multiplicative group $\mathbb{R}_+^\times$ of positive real numbers can be realized as the fundamental group $\mathcal{F}(A)$ of a separable simple unital C^* -algebra A with unique trace. Furthermore for any fixed countable subgroup G of $\mathbb{R}_+^\times$, there exist uncountably many mutually nonisomorphic such algebras A with $G = \mathcal{F}(A)$.

1. INTRODUCTION

Let M be a factor of type II_1 with a normalized trace τ . Murray and von Neumann introduced the fundamental group $\mathcal{F}(M)$ of M in [13]. The fundamental group $\mathcal{F}(M)$ of M is a subgroup of the multiplicative group $\mathbb{R}_+^\times$ of positive real numbers. They showed that if M is hyperfinite, then $\mathcal{F}(M) = \mathbb{R}_+^\times$. In our previous paper [14], we introduced the fundamental group $\mathcal{F}(A)$ of a simple unital C^* -algebra A with a unique normalized trace τ based on the computation of Picard groups by Kodaka [8], [9], [10]. We compute the fundamental groups of several nuclear or nonnuclear C^* -algebras. K -theoretical obstruction enable us to compute the fundamental group easily.

There has been many works on the computation of fundamental groups of the factors of type II_1 . Voiculescu [22] showed that the fundamental group $\mathcal{F}(L(\mathbb{F}_\infty))$ of the group factor $L(\mathbb{F}_\infty)$ of the free group $\mathbb{F}_\infty$ contains the positive rationals and Radulescu proved that $\mathcal{F}(L(\mathbb{F}_\infty)) = \mathbb{R}_+^\times$ in [19]. Connes [3] showed that if G is an countable ICC group with property (T), then $\mathcal{F}(L(G))$ is a countable group. Recently, Popa showed that any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group of some factor of type II_1 with separable predual in [17]. Furthermore Popa and Vaes [18] exhibited a large family $\mathcal{S}$ of subgroups of $\mathbb{R}_+^\times$, containing $\mathbb{R}_+^\times$ itself, all of its countable subgroups, as well as uncountable subgroups with any Hausdorff dimension in $(0, 1)$, such that for each $G \in \mathcal{S}$ there exist many free ergodic measure preserving actions of $\mathbb{F}_\infty$ for which the associated II_1 factor M has fundamental group equal to G .

In this paper we show that any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group of a separable simple unital C^* -algebra with unique trace. Furthermore for any fixed countable subgroup G of $\mathbb{R}_+^\times$, there exist uncountably many mutually nonisomorphic such algebras A with $G = \mathcal{F}(A)$. We apply a method of Blackadar [1] and Phillips [16] to the type II_1 factors of Popa [17]. Our new examples are nonnuclear.

On the other hand, for an additive subgroup E of $\mathbb{R}$ containing 1, we define the positive inner multiplier group $IM_+(E)$ of E by

$$IM_+(E) = \{t \in \mathbb{R}_+^\times \mid t \in E, t^{-1} \in E, \text{ and } tE = E\}.$$

Then we have $\mathcal{F}(A) \subset IM_+(\tau_*(K_0(A)))$. Almost all examples provided in [14] satisfy $\mathcal{F}(A) = IM_+(\tau_*(K_0(A)))$. We should note that not all countable subgroups of $\mathbb{R}_+^\times$ arise as $IM_+(E)$. For example, $\{9^n \in \mathbb{R}_+^\times \mid n \in \mathbb{Z}\}$ does not arise as $IM_+(E)$ for any additive subgroup E of $\mathbb{R}$ containing 1. Therefore if the fundamental group of a C^* -algebra A is equal to $\{9^n \in \mathbb{R}_+^\times \mid n \in \mathbb{Z}\}$ and A is in a classifiable class by the Elliott invariant, then $\tau_* : K_0(A) \rightarrow \tau_*(K_0(A))$ cannot be an order isomorphism. Matui informed us that there exists such an AF-algebra.

2. HILBERT C^* -MODULES AND PICARD GROUPS

We recall some definitions and notations in [14]. Let A be a simple unital C^* -algebra with a unique normalized trace τ and $\mathcal{X}$ a right Hilbert A -module. (See [11], [12] for the basic facts on Hilbert modules.) We denote by $L_A(\mathcal{X})$ the algebra of the adjointable operators on $\mathcal{X}$. For $\xi, \eta \in \mathcal{X}$, a "rank one operator" $\Theta_{\xi, \eta}$ is defined by $\Theta_{\xi, \eta}(\zeta) = \xi \langle \eta, \zeta \rangle_A$ for $\zeta \in \mathcal{X}$. We denote by $K_A(\mathcal{X})$ the closure of the linear span of "rank one operators" $\Theta_{\xi, \eta}$. We call a finite set $\{\xi_i\}_{i=1}^n \subseteq \mathcal{X}$ a *finite basis* of $\mathcal{X}$ if $\eta = \sum_{i=1}^n \xi_i \langle \xi_i, \eta \rangle_A$ for any $\eta \in \mathcal{X}$, see [7], [23]. It is also called a frame as in [6]. If A is unital and there exists a finite basis for $\mathcal{X}$, then $L_A(\mathcal{X}) = K_A(\mathcal{X})$. Let $\mathcal{H}(A)$ denote the set of isomorphic classes $[\mathcal{X}]$ of right Hilbert A -modules $\mathcal{X}$ with finite basis.

Let B be a C^* -algebra. An A - B -equivalence bimodule is an A - B -bimodule $\mathcal{F}$ which is simultaneously a full left Hilbert A -module under a left A -valued inner product ${}_A\langle \cdot, \cdot \rangle$ and a full right Hilbert B -module under a right B -valued inner product $\langle \cdot, \cdot \rangle_B$, satisfying ${}_A\langle \xi, \eta \rangle \zeta = \xi \langle \eta, \zeta \rangle_B$ for any $\xi, \eta, \zeta \in \mathcal{F}$. We say that A is *Morita equivalent* to B if there exists an A - B -equivalence bimodule. The dual module $\mathcal{F}^*$ of an A - B -equivalence bimodule $\mathcal{F}$ is a set $\{\xi^*; \xi \in \mathcal{F}\}$ with the operations such that $\xi^* + \eta^* = (\xi + \eta)^*$, $\lambda \xi^* = (\overline{\lambda} \xi)^*$, $b \xi^* a = (a^* \xi b^*)^*$, ${}_B\langle \xi^*, \eta^* \rangle = \langle \eta, \xi \rangle_B$ and $\langle \xi^*, \eta^* \rangle_A = {}_A\langle \eta, \xi \rangle$. Then $\mathcal{F}^*$ is a B - A -equivalence bimodule. We refer the reader to [20], [21] for the basic facts on equivalence bimodules and Morita equivalence.

We review elementary facts on the Picard groups of C^* -algebras introduced by Brown, Green and Rieffel in [2]. For A - A -equivalence bimodules $\mathcal{E}_1$ and $\mathcal{E}_2$, we say that $\mathcal{E}_1$ is isomorphic to $\mathcal{E}_2$ as an equivalence bimodule if there exists a $\mathbb{C}$ -linear one-to-one map Φ of $\mathcal{E}_1$ onto $\mathcal{E}_2$ with the properties such that $\Phi(a\xi b) = a\Phi(\xi)b$, ${}_A\langle \Phi(\xi), \Phi(\eta) \rangle = {}_A\langle \xi, \eta \rangle$ and $\langle \Phi(\xi), \Phi(\eta) \rangle_A = \langle \xi, \eta \rangle_A$ for $a, b \in A$, $\xi, \eta \in \mathcal{E}_1$. The set of isomorphic classes $[\mathcal{E}]$ of the A - A -equivalence bimodules $\mathcal{E}$ forms a group under the product defined by $[\mathcal{E}_1][\mathcal{E}_2] = [\mathcal{E}_1 \otimes_A \mathcal{E}_2]$. We call it the *Picard group* of A and denote it by $\text{Pic}(A)$. The identity of $\text{Pic}(A)$ is given by the A - A -bimodule $\mathcal{E} := A$ with ${}_A\langle a_1, a_2 \rangle = a_1 a_2^*$ and $\langle a_1, a_2 \rangle_A = a_1^* a_2$ for $a_1, a_2 \in A$. The inverse element of $[\mathcal{E}]$ in the Picard group of A is the dual module $[\mathcal{E}^*]$. Let α be an automorphism of A , and let $\mathcal{E}_\alpha^A = A$ with the obvious left A -action and the obvious A -valued inner product. We define the right A -action on $\mathcal{E}_\alpha^A$ by $\xi \cdot a = \xi \alpha(a)$ for any $\xi \in \mathcal{E}_\alpha^A$ and $a \in A$, and the right A -valued inner product by $\langle \xi, \eta \rangle_A = \alpha^{-1}(\xi^* \eta)$ for

any $\xi, \eta \in \mathcal{E}_\alpha^A$. Then $\mathcal{E}_\alpha^A$ is an A - A -equivalence bimodule. For $\alpha, \beta \in \text{Aut}(A)$, $\mathcal{E}_\alpha^A$ is isomorphic to $\mathcal{E}_\beta^A$ if and only if there exists a unitary $u \in A$ such that $\alpha = \text{ad } u \circ \beta$. Moreover, $\mathcal{E}_\alpha^A \otimes \mathcal{E}_\beta^A$ is isomorphic to $\mathcal{E}_{\alpha \circ \beta}^A$. Hence we obtain an homomorphism ρ_A of $\text{Out}(A)$ to $\text{Pic}(A)$. An A - B -equivalence bimodule $\mathcal{F}$ induces an isomorphism Ψ of $\text{Pic}(A)$ to $\text{Pic}(B)$ by $\Psi([\mathcal{E}]) = [\mathcal{F}^* \otimes \mathcal{E} \otimes \mathcal{F}]$ for $[\mathcal{E}] \in \text{Pic}(A)$. Therefore if A is Morita equivalent to B , then $\text{Pic}(A)$ is isomorphic to $\text{Pic}(B)$. Since A is unital, any A - A -equivalence bimodule is a finitely generated projective A -module as a right module with a finite basis $\{\xi_i\}_{i=1}^n$. Put $p = (\langle \xi_i, \xi_j \rangle_A)_{ij} \in M_n(A)$. Then p is a projection and $\mathcal{E}$ is isomorphic to pA^n as a right Hilbert A -module with an isomorphism of A to $pM_n(A)p$ as a C^* -algebra.

Define a map $\hat{T}_A : \mathcal{H}(A) \rightarrow \mathbb{R}_+$ by $\hat{T}_A([\mathcal{X}]) = \sum_{i=1}^n \tau(\langle \xi_i, \xi_i \rangle_A)$, where $\{\xi_i\}_{i=1}^n$ is a finite basis of $\mathcal{X}$. Then $\hat{T}_A([\mathcal{X}])$ does not depend on the choice of basis and $\hat{T}_A$ is well-defined. We can define a map T_A of $\text{Pic}(A)$ to $\mathbb{R}_+$ by the same way of $\hat{T}_A$. We showed that T_A is a multiplicative map and $T_A(\mathcal{E}_{id}^A) = 1$ in [14]. Moreover, we can show the following proposition by a similar argument in the proof of Proposition 2.1 in [14].

Proposition 2.1. Let A and B be simple unital C^* -algebras with unique trace. Assume that $\mathcal{F}$ is an A - B -equivalence bimodule and $\mathcal{X}$ is a right Hilbert A -module. Then

$$\hat{T}_B([\mathcal{X} \otimes \mathcal{F}]) = \hat{T}_A([\mathcal{X}])\hat{T}_B([\mathcal{F}]).$$

We denote by Tr the usual unnormalized trace on $M_n(\mathbb{C})$. Put

$$\mathcal{F}(A) := \{\tau \otimes Tr(p) \in \mathbb{R}_+^\times \mid p \text{ is a projection in } M_n(A) \text{ such that } pM_n(A)p \cong A\}.$$

Then $\mathcal{F}(A)$ is equal to the image of T_A and a multiplicative subgroup of $\mathbb{R}_+^\times$ by Theorem 3.1 in [14]. We call $\mathcal{F}(A)$ the *fundamental group* of A . If A is separable, then $\mathcal{F}(A)$ is countable. We shall show that the fundamental group is a Morita equivalence invariant for simple unital C^* -algebras with unique trace.

Proposition 2.2. Let A and B be simple unital C^* -algebras with unique trace. If A is Morita equivalent to B , then $\mathcal{F}(A) = \mathcal{F}(B)$.

Proof. By assumption, there exists an A - B -equivalence bimodule $\mathcal{F}$, and $\mathcal{F}$ induces an isomorphism Ψ of $\text{Pic}(A)$ to $\text{Pic}(B)$ such that $\Psi([\mathcal{E}]) = [\mathcal{F}^* \otimes \mathcal{E} \otimes \mathcal{F}]$ for $[\mathcal{E}] \in \text{Pic}(A)$. Since $\mathcal{F}^* \otimes \mathcal{F}$ is isomorphic to $\mathcal{E}_{id}^B$, Proposition 2.1 implies

$$\hat{T}_A([\mathcal{F}^*])\hat{T}_B([\mathcal{F}]) = T_B([\mathcal{F}^* \otimes \mathcal{F}]) = 1.$$

For $[\mathcal{E}] \in \text{Pic}(A)$,

$$T_B([\mathcal{F}^* \otimes \mathcal{E} \otimes \mathcal{F}]) = \hat{T}_A([\mathcal{F}^*])\hat{T}_B([\mathcal{E} \otimes \mathcal{F}]) = \hat{T}_A([\mathcal{F}^*])T_A([\mathcal{E}])\hat{T}_B([\mathcal{F}])$$

by Proposition 2.1. Therefore $T_B([\Psi(\mathcal{E})]) = T_A([\mathcal{E}])$ and $\mathcal{F}(A) = \mathcal{F}(B)$. $\square$

3. NEW EXAMPLES

An idea of our construction comes from the following results of Blackadar, Proposition 2.2 of [1] and Phillips, Lemma 2.2 of [16].

Lemma 3.1 ([1](Blackadar)). Let M be a simple C^* -algebra, and let $A \subset M$ be a separable C^* -subalgebra. Then there exists a simple separable C^* -subalgebra B with $A \subset B \subset M$.

Lemma 3.2 ([16](Phillips)). Let M be a unital C^* -algebra, and let $A \subset M$ be a separable C^* -subalgebra. Then there exists a separable C^* -subalgebra B with $A \subset B \subset M$ such that every tracial state on B is the restriction of a tracial state on M .

The following lemma is just a combination of the two results above.

Lemma 3.3. Let M be a simple C^* -algebra with unique trace $\hat{\tau}$, and let $A \subset M$ be a separable C^* -subalgebra. Then there exists a simple separable C^* -subalgebra B with $A \subset B \subset M$ such that B has a unique trace τ that is a restriction of $\hat{\tau}$.

Theorem 3.4. Let G be a countable subgroup of $\mathbb{R}_+^\times$. Then there exist uncountably many mutually nonisomorphic separable simple nonnuclear unital C^* -algebras A with unique trace such that the fundamental group $\mathcal{F}(A) = G$.

Proof. First we shall show that there exists a separable simple unital C^* -algebra A with unique trace such that $\mathcal{F}(A) = G$. There exists a type II_1 factor M with separable predual such that $\mathcal{F}(M) = G$, which is constructed by Popa [17]. Let $S_1 \subset M$ be a countable subset that is weak operator dense in M . We denote by $\hat{\tau}$ the unique trace of M . We enumerate the countable semigroup $G \cap (0, 1]$ by $\{t_m : m \in \mathbb{N}\}$. Since $\mathcal{F}(M) = G$ and M is a factor of type II_1 , for any $m \in \mathbb{N}$ there exist a projection p_m in M such that $\hat{\tau}(p_m) = t_m$ and an isomorphism ϕ_m of M onto $p_m M p_m$. Define $B_0 \subset M$ be the unital C^* -subalgebra of M generated by S_1 and $\{p_m : m \in \mathbb{N}\}$. By Lemma 3.3, there exists a separable simple unital C^* -algebra A_0 with a unique trace τ_0 such that $B_0 \subset A_0 \subset M$. Let $B_1 \subset M$ be the C^* -subalgebra of M generated by A_0 , $\cup_{m \in \mathbb{N}} \phi_m(A_0)$ and $\cup_{m \in \mathbb{N}} \phi_m^{-1}(p_m A_0 p_m)$. By the same way, there exists a separable simple unital C^* -algebra A_1 with a unique trace τ_1 such that $B_1 \subset A_1 \subset M$. We construct inductively C^* -algebras $B_n \subset A_n \subset M$ as follows: Let $B_n \subset M$ be the C^* -subalgebra of M generated by A_{n-1} , $\cup_{m \in \mathbb{N}} \phi_m(A_{n-1})$ and $\cup_{m \in \mathbb{N}} \phi_m^{-1}(p_m A_{n-1} p_m)$. By Lemma 3.3, there exists a separable simple unital C^* -algebra A_n with a unique trace τ_n such that $B_n \subset A_n \subset M$. Then we have

$$B_0 \subset A_0 \subset B_1 \subset A_1 \subset \dots B_n \subset A_n \subset \dots \subset M,$$

and $\phi_m(A_{n-1}) \subset p_m A_n p_m$ and $\phi_m^{-1}(p_m A_{n-1} p_m) \subset A_n$ for any $m \in \mathbb{N}$. Set $A = \overline{\cup_{n=0}^\infty A_n}$. Then A is a separable simple unital C^* -algebra A with a unique trace τ . By the construction, $\phi_m(A) = p_m A p_m$ for any $m \in \mathbb{N}$. Hence $G \subset \mathcal{F}(A)$. Since $\pi_\tau(A)''$ is isomorphic to M ,

$$\mathcal{F}(A) \subset \mathcal{F}(\pi_\tau(A)'') = \mathcal{F}(M) = G$$

by Proposition 3.29 of [14]. Thus $\mathcal{F}(A) = G$. Moreover A is not nuclear, because A is weak operator dense in a factor M that is not hyperfinite.

Next we shall show that there exist uncountably many mutually nonisomorphic such examples. Let E be a countable additive subgroup of $\mathbb{R}$. We enumerate by $\{r_m : m \in \mathbb{N}\}$ the positive elements of E . Since M is a factor of

type II_1 , for any $m \in \mathbb{N}$ there exist a natural number k and a projection $q_m \in M_k(M)$ such that $\hat{\tau} \otimes \text{Tr}(q_m) = r_m$. Define $S_2 \subset M$ to be the union of the matrix elements of q_m for running $m \in \mathbb{N}$. Let C_0 the C^* -subalgebra of M generated by S_2 and A . By a similar argument as the first paragraph, we can construct a separable simple unital C^* -algebra C with unique trace such that $\mathcal{F}(C) = G$ and $C_0 \subset C \subset M$. Then it is clear that E is contained in $\tau_*(K_0(C))$. Since no countable union of countable subgroups of $\mathbb{R}$ can contain all countable subgroups of $\mathbb{R}$, we can construct uncountably many mutually nonisomorphic examples by the choice of E . $\square$

Remark 3.5. In fact, we show that there exist uncountably many Morita inequivalent separable simple nonnuclear unital C^* -algebras A with unique trace such that the fundamental group $\mathcal{F}(A) = G$ in the proof above.

Remark 3.6. We can choose a C^* -algebra A in the theorem above so that A has stable rank one and real rank zero and $\tau_* : K_0(A) \rightarrow \tau_*(K_0(A))$ is an order isomorphism by using Lemma 2.3, Lemma 2.4 and Lemma 2.5 of [16]. Then we have the following exact sequence by Proposition 3.26 of [14]:

$$1 \longrightarrow \text{Out}(A) \xrightarrow{\rho_A} \text{Pic}(A) \xrightarrow{T} \mathcal{F}(A) \longrightarrow 1.$$

Remark 3.7. We do not know whether any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group of a separable unital simple *nuclear* C^* -algebra with unique trace.

Lemma 3.8. Let M_1 and M_2 be factors of type II_1 , and let $A_0 \subset M_1$ and $B_0 \subset M_2$ be separable C^* -subalgebras. Then there exist separable simple unital C^* -algebras A and B with the unique traces τ_A and τ_B such that $A_0 \subset A \subset M_1$, $B_0 \subset B \subset M_2$ and $(\tau_A)_*(K_0(A)) = (\tau_B)_*(K_0(B))$.

Proof. Let τ_1 be the unique trace on M_1 and τ_2 the unique trace on M_2 . Since A_0 and B_0 are separable C^* -algebras, $(\tau_1|_{A_0})_*(K_0(A_0))$ and $(\tau_2|_{B_0})_*(K_0(B_0))$ are countable groups. We enumerate the positive elements of $(\tau_1|_{A_0})_*(K_0(A_0))$ by $\{t_m : m \in \mathbb{N}\}$ and the positive elements of $(\tau_2|_{B_0})_*(K_0(B_0))$ by $\{r_m : m \in \mathbb{N}\}$. Since M_1 and M_2 are factors of type II_1 , for any $m \in \mathbb{N}$ there exist a natural number k and projections $p_m \in M_k(M_1)$ and $q_m \in M_k(M_2)$ such that $\tau_1 \otimes \text{Tr}(p_m) = r_m$ and $\tau_2 \otimes \text{Tr}(q_m) = t_m$. Put $S_1 \subset M_1$ (resp. $S_2 \subset M_2$) to be the union of the matrix elements of p_m (resp. q_m) for running $m \in \mathbb{N}$. Define $C_1 \subset M_1$ (resp. $D_1 \subset M_2$) be the unital C^* -subalgebra of M_1 (resp. M_2) generated by A_0 and S_1 (resp. B_0 and S_2). By Lemma 3.3, there exist separable simple unital C^* -algebras A_1 and B_1 with a unique trace such that $C_1 \subset A_1 \subset M_1$ and $D_1 \subset B_1 \subset M_2$. Then we have $(\tau_1|_{A_0})_*(K_0(A_0)) \subset (\tau_2|_{B_1})_*(K_0(B_1))$ and $(\tau_2|_{B_0})_*(K_0(B_0)) \subset (\tau_1|_{A_1})_*(K_0(A_1))$. In a similar way, we construct inductively simple separable unital C^* -algebras $A_n \subset M_1$ and $B_n \subset M_2$ with unique trace such that $(\tau_1|_{A_{n-1}})_*(K_0(A_{n-1})) \subset (\tau_2|_{B_n})_*(K_0(B_n))$ and $(\tau_2|_{B_{n-1}})_*(K_0(B_{n-1})) \subset (\tau_1|_{A_n})_*(K_0(A_n))$. Set $A = \overline{\bigcup_{n=1}^\infty A_n}$ and $B = \overline{\bigcup_{n=1}^\infty B_n}$. Then A and B are separable simple unital C^* -algebras with unique trace. We denote by τ_A the unique trace on A and by τ_B the unique traces on B . By the construction, $(\tau_A)_*(K_0(A)) = (\tau_B)_*(K_0(B))$. $\square$

We denote by $\text{Ell}(A)$ the Elliott invariant $(K_0(A), K_0(A)_+, [1]_0, K_1(A))$.

Corollary 3.9. For any countable subgroups G_1 and G_2 of $\mathbb{R}_+^\times$, there exist separable simple nonnuclear unital C^* -algebras A and B with unique trace such that $\text{Ell}(A) \cong \text{Ell}(B)$, $\mathcal{F}(A) = G_1$ and $\mathcal{F}(B) = G_2$.

Proof. The proof of Theorem 3.4, Lemma 3.8 and Lemma 2.5 of [16] implies that there exist separable simple nonnuclear unital C^* -algebras A and B with the unique traces τ_A and τ_B such that $(\tau_A)_* : K_0(A) \rightarrow (\tau_A)_*(K_0(A))$ and $(\tau_B)_* : K_0(B) \rightarrow (\tau_B)_*(K_0(B))$ are order isomorphisms, $\mathcal{F}(A) = G_1$, $\mathcal{F}(B) = G_2$, $K_1(A) = K_1(B) = 0$ and $(\tau_A)_*(K_0(A)) = (\tau_B)_*(K_0(B))$. Since $(\tau_A)_*$ and $(\tau_B)_*$ are order isomorphisms and $(\tau_A)_*(K_0(A)) = (\tau_B)_*(K_0(B))$, we see that $\text{Ell}(A) \cong \text{Ell}(B)$. $\square$

For a positive number λ , let $G_\lambda = \{\lambda^n \in \mathbb{R}_+^\times \mid n \in \mathbb{Z}\}$ be the multiplicative subgroup of $\mathbb{R}_+^\times$ generated by λ . In the below we shall consider whether G_λ can be realized as the fundamental group of a nuclear C^* -algebra.

Proposition 3.10. Let λ be a prime number or a positive transcendental number. Then there exists a simple AF -algebra A with unique trace such that $\mathcal{F}(A) = G_\lambda$.

Proof. Let λ be a prime number. Consider a UHF-algebra $A = M_\lambda^\infty$. Then $\mathcal{F}(A) = G_\lambda$ as in Example 3.11 of [14]. Next we assume that λ is a positive transcendental number. Let R_λ be the unital subring of $\mathbb{R}$ generated by λ . Then the set $(R_\lambda)_+^\times$ of positive invertible elements in R_λ is equal to G_λ . The proof of Theorem 3.14 of [14] shows that there exists a simple unital AF -algebra A with unique trace such that $\mathcal{F}(A) = G_\lambda$. $\square$

Let $\mathcal{O}$ be an order of a real quadratic field or a real cubic field with one real embedding. Then $\mathcal{O}_+^\times = G_\lambda$ is singly generated and the generator $\lambda > 1$ is called the fundamental unit of $\mathcal{O}$ by Dirichlet's unit theorem. We refer the reader to [15] for details. The proof of Theorem 3.14 of [14] implies the following proposition.

Proposition 3.11. Let λ be a fundamental unit of an order of a real quadratic field or a cubic field with one real embedding. Then there exists a simple AF -algebra A with unique trace such that $\mathcal{F}(A) = G_\lambda$.

Note that if p is a prime number and $n \geq 2$, then the subgroup G_λ of $R_+^\times$ generated by $\lambda = p^n$ can not be the positive inner multiplier group $IM_+(E)$ for any additive subgroup E of $\mathbb{R}$ containing 1. In fact, on the contrary, suppose that $G_\lambda = IM_+(E)$ for some E . Then there exists a unital subring R of $\mathbb{R}$ such that $G_\lambda = R_+^\times$ by Lemma 3.6 of [14]. Then $\frac{1}{p} = \frac{1}{\lambda} + \dots + \frac{1}{\lambda} \in R_+^\times$. This contradicts that $\frac{1}{p} \notin G_\lambda$. However, we have another construction.

Example 3.12. For $\lambda = 3^2 = 9$, Matui shows us the following example: Let A be an AF -algebra such that

$$K_0(A) = \{(\frac{b}{9^a}, c) \in \mathbb{R} \times \mathbb{Z} \mid a, b, c \in \mathbb{Z}, b \equiv c \pmod{8}\},$$

$$K_0(A)_+ = \{(\frac{b}{9^a}, c) \in K_0(A) : \frac{b}{9^a} > 0\} \cup \{(0, 0)\} \quad \text{and} \quad [1_A]_0 = (1, 1).$$

Then

$$\mathcal{F}(A) = G_9 := \{9^n \in \mathbb{R}_+^\times \mid n \in \mathbb{Z}\}$$

Moreover $\tau_* : K_0(A) \rightarrow \tau_*(K_0(A))$ is not an order isomorphism and $\mathcal{F}(A) \neq IM_+(\tau_*(K_0(A)))$.

Furthermore Katsura suggests us the following examples: Let $\lambda = p^n$ for a prime number p and a natural number $n \geq 2$. Then there exists a simple AF -algebra A with unique trace such that $\mathcal{F}(A) = G_\lambda$.

First consider the case that $\lambda \geq 8$. Define

$$E = \{(\frac{b}{p^{na}}, c) \in \mathbb{R} \times \mathbb{Z} \mid a, b, c \in \mathbb{Z}, b \equiv c \pmod{p^n - 1}\}$$

$$E_+ = \{(\frac{b}{p^{na}}, c) \in E : \frac{b}{p^{na}} > 0\} \cup \{(0, 0)\} \quad \text{and} \quad [u]_0 = (1, 1).$$

Then there exists a simple AF -algebra A such that $(K_0(A), K_0(A)_+, [1_A]_0) = (E, E_+, u)$ by [4]. The classification theorem of [5] and some computation yield that $\mathcal{F}(A) = G_\lambda$.

Next consider the case that $\lambda = 2^2 = 4$. Let

$$E = \{(\frac{b}{16^a}, c) \in \mathbb{R} \times \mathbb{Z} \mid a, b, c \in \mathbb{Z}, b \equiv c \pmod{5}\}$$

$$E_+ = \{(\frac{b}{16^a}, c) \in E : \frac{b}{16^a} > 0\} \cup \{(0, 0)\} \quad \text{and} \quad [u]_0 = (1, 1).$$

Consider a simple AF -algebra A such that $(K_0(A), K_0(A)_+, [1_A]_0) = (E, E_+, u)$. Then $\mathcal{F}(A) = G_4$.

REFERENCES

- [1] B. Blackadar, *Weak expectations and nuclear C^* -algebras*, Indiana Univ. Math. J. **27** (1978), 1021-1026.
- [2] L. G. Brown, P. Green and M. A. Rieffel, *Stable isomorphism and strong Morita equivalence of C^* -algebras*, Pacific J. Math. **71** (1977), 349-363.
- [3] A. Connes, *A factor of type II_1 with countable fundamental group*, J. Operator Theory **4** (1980), 151-153.
- [4] E. Effros, D. Handelman and C. L. Shen, *Dimension groups and their affine representations*, Amer. J. Math. **102** (1980), no. 2, 385-407.
- [5] G. A. Elliott, *On the classification of inductive limits of sequences of semisimple finite-dimensional algebras*, J. Algebra **38** (1976), 29-44.
- [6] M. Frank and D. Larson, *A module frame concept for Hilbert C^* -modules*, Contemporary Mathematics **247** (1999), 207-233.
- [7] T. Kajiwara and Y. Watatani, *Jones index theory by Hilbert C^* -bimodules and K -theory*, Trans. Amer. Math. Soc. **352** (2000), 3429-3472.
- [8] K. Kodaka, *Full projections, equivalence bimodules and automorphisms of stable algebras of unital C^* -algebras*, J. Operator Theory, **37** (1997), 357-369.
- [9] K. Kodaka, *Picard groups of irrational rotation C^* -algebras*, J. London Math. Soc. (2) **56** (1997), 179-188.
- [10] K. Kodaka, *Projections inducing automorphisms of stable UHF-algebras*, Glasg. Math. J. **41** (1999), no. 3, 345-354.
- [11] E. C. Lance, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series, **210**, Cambridge University Press, Cambridge, 1995.
- [12] V. M. Manuilov and E. V. Troitsky, *Hilbert C^* -Modules*, Translations of Mathematical Monographs, **226**, American Mathematical Society, Providence, RI, 2005.
- [13] F. Murray and J. von Neumann, *On rings of operators IV*, Ann. Math. **44**, (1943), 716-808.
- [14] N. Nawata and Y. Watatani, *Fundamental group of simple C^* -algebras with unique trace*, to appear in Adv. Math.
- [15] J. Neukirch, *Algebraic Number Theory*, Grundlehren Math. Wiss. vol. **322**, Springer-Verlag, New York, 1999.

- [16] N.C. Phillips, *A Simple separable C^* -algebra not isomorphic to its opposite algebra*, Proc. Amer. Math. Soc. **132** (10) (2004), 2997-3005.
- [17] S. Popa, *Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups, I*, Invent. Math. **165** (2006), 369-408.
- [18] S. Popa and S. Vaes, *Actions of $\mathbb{F}_\infty$ whose II_1 factors and orbit equivalence relations have prescribed fundamental group*, J. Amer. Math. Soc. **23** (2010), 383-403.
- [19] F. Radulescu, *The fundamental group of the von Neumann algebra of a free group with infinitely many generators is $\mathbb{R}_+$* , J. Amer. Math. Soc. **5** (1992), 517-532.
- [20] I. Raeburn and D. P. Williams, *Morita Equivalence and Continuous-Trace C^* -Algebras*, Mathematical Surveys and Monographs, **60**, American Mathematical Society, Providence, RI, 1998.
- [21] M. A. Rieffel, *Morita equivalence for operator algebras*, Operator algebras and applications, Part I (Kingston, Ont., 1980), pp. 285-298, Proc. Sympos. Pure Math., **38**, Amer. Math. Soc., Providence, R.I., 1982.
- [22] D. Voiculescu, *Circular and semicircular systems and free product factors*, in Operator algebras, unitary representations, enveloping algebras, and invariant theory, Progr. Math. 92, Birkhäuser, Boston, 1990, 45-60.
- [23] Y. Watatani, *Index for C^* -subalgebras*, Memoir AMS **424** (1990).

(Norio Nawata) GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY, MOTOOKA, FUKUOKA, 819-0395, JAPAN

(Yasuo Watatani) DEPARTMENT OF MATHEMATICAL SCIENCES, KYUSHU UNIVERSITY, MOTOOKA, FUKUOKA, 819-0395, JAPAN